

Neurodegenerative Diseases in Welders and Other Workers Exposed to High Levels of Magnetic Fields

Niclas Håkansson ^{*}†; Per Gustavsson [‡]§; Christoffer Johansen ¶; Birgitta Floderus †

Background:

Previous work has suggested an increase in risk of amyotrophic lateral sclerosis (ALS) and Alzheimer's disease among workers exposed to extremely low-frequency magnetic fields (ELF-MF). We evaluated the relation between ELF-MF from occupational exposures and mortality from neurodegenerative diseases.

Methods:

The study was based on a cohort of Swedish engineering industry workers, comprising 537,692 men and 180,529 women. The cohort was matched against the 3 most recent censuses and The Causes of Death Registry. Levels of ELF-MF exposure were obtained by linking occupation according to the censuses to a job exposure matrix. We used 4 levels of exposure and considered both the primary and contributing causes of death, 1985-96.

Results:

The risk of Alzheimer's disease as primary or contributing cause of death increased with increasing exposure to ELF-MF among both men and women, with a relative risk (RR) of 4.0 and a 95% confidence interval (95% CI) of 1.4-11.7 in the highest exposure group for both sexes combined. There was a RR of 2.2 (95% CI: 1.0-4.7) for ALS in the highest exposure group with the suggestion of an exposure-response relationship. No evidence of increased risk was seen for Parkinson's disease or multiple sclerosis.

Conclusions:

The findings support previous observations of an increased risk of Alzheimer's disease and ALS among employees occupationally exposed to ELF-MF. Further studies based on morbidity data are warranted.

Key Words Alzheimer's disease; Amyotrophic Lateral Sclerosis; Electromagnetic Fields; Neurodegenerative Disease; Occupation

From the ^{*}Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden, [†]National Institute for Working Life, Solna, Sweden, [‡]Department of Occupational and Environmental Health, Stockholm Public Health Center, Stockholm, Sweden, [§]Division of Occupational Health, Department of Public Health Sciences, Karolinska Institutet, Stockholm, Sweden, [¶]Institute of Cancer Epidemiology, Danish Cancer Society, Copenhagen, Denmark

This work was supported by grant from ELFORSK, Sweden.

C

Address correspondence to: Niclas Håkansson, Institute of Environmental Medicine, Karolinska Institutet, Box 210, S-171 77 Stockholm, Sweden. Tel: +46 8 728 70 80, Fax: +46 8 30 45 71. E-mail: niclas.hakansson@imm.ki.se.

Submitted 12 March 2002;

final version accepted 12 November 2002.

Epidemiology 2003; 14(4):420-426
Copyright © 2003 Lippincott Williams & Wilkins
All rights reserved

O

N

A

C

E

M